

AD A 056906

DDC FILE COPY



LEVEL II

(11)

AD

Report 2247

ENERGY DEVELOPMENT FROM ELEMENTAL  
TRANSmutations IN BIOLOGICAL SYSTEMS

by  
S. Goldfein

May 1978

DDC  
APR 2 1978  
F

Approved for public release; distribution unlimited.

U.S. ARMY MOBILITY EQUIPMENT  
RESEARCH AND DEVELOPMENT COMMAND  
FORT BELVOIR, VIRGINIA

78 07 27 054

Destroy this report when it is no longer needed.  
Do not return it to the originator.

The citation in this report of trade names of commercially  
available products does not constitute official endorsement  
or approval of the use of such products.

New Energy Times

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2247	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ENERGY DEVELOPMENT FROM ELEMENTAL TRANSMUTATIONS IN BIOLOGICAL SYSTEMS		5. TYPE OF REPORT & PERIOD COVERED Final <sup>repts</sup> Dec 77 - April 78
7. AUTHOR(s) S/Goldfein		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Army Mobility Equipment Research and Development Command, DRDME-VL Fort Belvoir, VA 22060		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS MIL B1901 PMMR-26
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE May 78
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 72 247 P.
		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract included in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) energy source biological transmutation of elements magnesium adenosine triphosphate molecular cyclotron		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of the study was to determine whether recent disclosures of elemental transmutations occurring in biological entities have revealed new possible sources of energy. The works of Kerran, Komaki, and others were surveyed; and it was concluded that, granted the existence of such transmutations (Na to Mg, K to Ca, and Mn to Fe), then a net surplus of energy was also produced. A proposed mechanism was described in which Mg adenosine triphosphate, located in the mitochondrion of the cell, played a double role as an energy producer. In addition to the widely accepted biochemical role of MgATP in which it produces energy as it disintegrates part (Continued)		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

443 160

over  
KC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

(Block 20 (Cont'd))

by part, MgATP can also be considered to be a cyclotron on a molecular scale. The MgATP when placed in layers one atop the other has all the attributes of a cyclotron in accordance with the requirements set forth by E. O. Lawrence, inventor of the cyclotron.

It was concluded that elemental transmutations were indeed occurring in life organisms and were probably accompanied by a net energy gain.

New Energy Times

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## PREFACE

The work described in this report was authorized and funded by the U.S. Army Mobility Equipment Research and Development Command, Material Technology Laboratory, under Project MTL 01901, PMMR-26.

Solomon Goldfein was the principal investigator for the effort. The work was under the direction of Emil J. York, Chief, Material Technology Laboratory.

Robert C. McMillan, Chief, Radiation Research Group, provided guidance with regard to matters relating to physics and nuclear physics aspects.

New Energy Times

ACCESSION FOR	
NTIS	<input checked="" type="checkbox"/>
DOC	<input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	<input type="checkbox"/>
BY	
DISTRIBUTION/ACCESSION NOTES	
DATE	BY
A	

## CONTENTS

Section	Title	Page
	PREFACE	iii
	ILLUSTRATIONS	v
I	INTRODUCTION	
	1. Subject	1
	2. Background	1
II	INVESTIGATION	
	3. Plan	2
	a. Net Weight Change	3
	b. Energy Required for Nuclear Reactions	4
	c. Probable Location of Energy Development	4
	d. Chemicals Involved	6
	e. Energy Production	6
	f. Ions Present in the Mitochondrion	6
	g. Current Flow	10
	h. Structure of MgATP	10
	4. Proposed Alternate Mechanism for Energy Production	10
	a. Oscillatory Electric Field	15
	b. Presence of Hydrogen Ions	15
	c. Auxiliary Control and Propulsion of H <sup>+</sup> by Dipoles	15
	d. Induction of Ring Current in Adenine	15
	e. Effect of Magnetic Field on Hydrogen Ion	15
	f. Circular Helical Path of Hydrogen Ion	16
III	DISCUSSION	
	5. General	16
IV	CONCLUSIONS	
	6. Conclusions	17
	REFERENCES	18

## ILLUSTRATIONS

Figure	Title	Page
1	Typical Cell Showing a Number of Mitochondria	5
2	Membrane of the Mitochondrion	7
3	Chemical Structure of Adenosine Triphosphate (ATP)	8
4	Chemical Structure of Adenosine Triphosphate Chelated with Magnesium Ion (MgATP)	9
5	Planar View of Two Molecules of MgATP with Six Dipoles	12
6	Planar View of Four Molecules of MgATP with Twelve Dipoles	13
7	Side View of Four Molecules of MgATP Showing Circular Spiral Path of H <sup>+</sup>	14

## ENERGY DEVELOPMENT FROM ELEMENTAL TRANSMUTATIONS IN BIOLOGICAL SYSTEMS

### I. INTRODUCTION

1. **Subject.** The object of this study was to determine whether recent disclosures of elemental transmutations occurring in biological entities have revealed new possible sources of energy.

2. **Background.** Two investigators, Kervran and Komaki,<sup>1, 2</sup> have been recently nominated for a joint Nobel prize for their work involving the experimental proof that elemental transmutations were occurring in life organisms. Elements which were definitely proven to have been transformed were sodium (to magnesium), potassium (to calcium), and manganese (to iron). Actually, observations have been made for almost 200 years that elemental mutations were occurring, but little credence was given to them because they resembled alchemy - a relic of the middle ages.

From Jueneman,<sup>3</sup> investigations on transmutations can be traced from 1799 to the present.

The French Chemist, Vauquelin, in 1799, observed a large quantity of lime in the daily excretion of hens. To determine where the lime was coming from, he fed a captive hen a diet of nothing but oats. He measured the amount of lime in the oats, fed the oats to the hen, and then measured the amount of lime in the excretion and the eggs of the hen. The lime had increased fivefold. Vauquelin realized that lime had been created but he did not know why.

In 1822, Prout, an English experimenter, defined the transmutation of elements by studying incubating chicken eggs. He observed increasing amounts of calcium carbonate inside the egg and determined that the compound did not come from the shell.

In 1831, Chouard experimented with germinating watercress seeds. He observed new minerals in the young plants that were not present in the original seeds.

In 1844, Vogel also worked with watercress seeds. He placed his seeds in a controlled-air environment and added a sulfur-free nutritive solution to the seeds. The plants that grew from the seeds were found to have more sulfur than the original seeds.

<sup>1</sup> C. L. Kerstan, *Prev. en Biologie de Transmutation a Faible Energy*, Malorinix, Paris (1975).

<sup>2</sup> H. Komaki, *Rev. de Pathologie Comparee*, 67, 213 (1967); 69, 29 (1969).

<sup>3</sup> F. B. Jueneman, "The Transmutation of Species," *Industrial Research*, 19, No. 13, 11 (Dec 1977).



Vogel's answer was that sulfur was not a simple element or that sulfur was introduced from sources unknown.

After Vogel, Lauwes and Gilbert observed the weight variation of ashes during vegetation. In particular, they found a variation in magnesium.

In 1875, Von Herzele also studied vegetating plants and discovered a weight increase in the ashes of the young plants.

Later, following the works of Vogel and Lauwes and Gilbert, Von Herzele considered the variation in the weight of magnesium. He concluded that a transmutation of elements had taken place.

In 1960, Baranger, of France, published his findings on variations of calcium and phosphorus in germinating seeds. He also concluded that a transmutation had taken place, but he did not determine the reason.

In the early sixties, Louis Kervran's discoveries on transmutations were published by a French scientific review. Kervran demonstrated that not only molecules but also atoms could be transformed, and he verified the transmutation of matter from atom to atom.<sup>4</sup>

Komaki reported that eight strains of microorganisms grown in K-deficient culture media increase the total K by converting Ca to K - a fission reaction. A theoretical basis for Kervran's result has been discussed by de Beauregard.<sup>5</sup>

Working at the Ecole Polytechnique in Paris, Baranger, who also headed the organic laboratory, determined that transmutations were indeed taking place. Yet, none of these latter-day alchemists could propose a theory for the mechanism of such an unlikely appearing event as biological transmutation.

Summarizing, disclosures have been made of transmutations in microorganisms, plants, and animals. Some have been fusion while others have been fission phenomena.

## II. INVESTIGATION

3. **Plan.** In the search for clues to the possibilities for energy development, calculations were made to determine whether there was a net weight loss during the


<sup>4</sup> V. B. Juergensen, "The Transmutation of Species," *Industrial Research*, 19, No. 13, 11 (Dec 1977).

<sup>5</sup> O. Costa de Beauregard, *Proc. 3rd Int. Cong. Psych.* (Tokyo), 168 (June 1967).


transmutations. This was followed by an examination of the most likely place for the event to occur and the presence there of the same elements disclosed to have undergone nuclear changes. A study then followed to determine if a mechanism was present which might cause such nuclear transmutations in accordance with presently accepted nuclear theory.

a. **Net Weight Change.** The study was limited to fusion reactions so that the difference in atomic number would be only one. These included Na, K, and Mn:


Sodium to magnesium where:






Potassium to calcium where:



Magnesium to iron where:



The atomic weight differences were:



Energy developed as a result of transmutation:

Conversion factor for atomic mass unit (amu) =  $931 \frac{\text{MeV}}{\text{amu}}$

Na $\longrightarrow$ Mg	0.077 825 19 (931)	(7.29 MeV)
K $\longrightarrow$ Ca	0.008 976 39 (931)	(+8.35 MeV)
Mn $\longrightarrow$ Fe	0.010 030 10 (931)	(+10.18 MeV)

b. **Energy Required for Nuclear Reactions.** No information has been discovered as to the energy required for these reactions to occur and, thus, whether there would be a net gain of energy. As far back as 1932, however, it was known that alpha particles having energies when ejected from radioactive elements of medium life of between 5 and 7 MeV could cause atomic disintegrations. Cockcroft and Walton built a tall, vertical tube capable of evacuation with a filament producing electrons at the top and a target of the element to be bombarded at the bottom. A low pressure of hydrogen was introduced and ionized by collisions with the electrons produced by the filament. The top of the tube could be raised to a high positive potential, up to 700,000 volts, the bottom being at earth. On reaching the bottom of the tube, the protons had a kinetic energy equal in electron volts to the potential in volts at the top of the tube. Using a lithium target, at  $45^\circ$  to the bombarding protons, disintegrations of some of the lithium atoms into pairs of helium nuclei ejected almost in opposite directions were observed with only 120,000 volts.<sup>6</sup>

The reaction is:



Another such pertinent reaction was:



c. **Probable Location of Energy Development.** The mitochondrion, a cylindrically shaped organelle, is universally recognized as the primary site for energy production in all life organisms regardless of whether they are one-celled bacteria, plants, or animals.<sup>7</sup> Some cells have as many as 7000 mitochondria (Figure 1). The

<sup>6</sup> J. Cockcroft and B. Lewis, *Proc. Roy. Soc. A*, 136, 619; 137, 229 (1932); 154, 246, 26 (1936).

<sup>7</sup> Henry R. Sgaller and Eugene H. Cordes, *Biological Chemistry* 17601, 609, 618 (1966).





Figure 1. Typical cell showing a number of mitochondria.

mitochondrion is divided into compartments in which various reactions take place (Figure 2).<sup>8</sup>

d. **Chemicals Involved.** Adenosine triphosphate (ATP) is now recognized as the molecule most involved in energy production in the mitochondrion (Figure 3). When complexed with  $Mg^{++}$ , it forms a cyclic ATP and proceeds through a series of reactions which produces energy (Figure 4).

e. **Energy Production.** The  $Mg^{++}$  is considered to be a catalyst for these reactions. One phosphate group after another splits off so that the resultant chelate becomes a diphosphate (ADP) and then a monophosphate (AMP). The phosphate groups hydrolyze, and the energy of hydrolysis for each reaction is 7,500 calories. The D-ribose then splits off and proceeds through a long glycolic decomposition cycle to yield further energy. The chelated, cyclic MgATP thus breaks up completely and is rebuilt through a series of chemical reactions. None of the decomposition and rebuilding cycles is as yet completely understood.

f. **Ions Present in the Mitochondrion.** Na, Mg, K, Ca, Mn, and Fe ions have been found to be present in the mitochondrion. These are the same ions which have been previously found to undergo nuclear transmutations by Kervran. In addition, another pair, Cu and Zn, differing by one proton (elements numbered 29 and 30) is also present.<sup>9</sup>  $H^+$  and  $OH^-$  ions have been produced and are maintained in separate compartments.<sup>10</sup> Conversion of Cu to Zn is also accompanied by a loss of mass and thus a production of energy:



<sup>8</sup> Henry R. McKer and Eugene H. Cordes, *Biological Chemistry*, 17 601, 609, 618 (1966).

<sup>9</sup> *Ibid.*

<sup>10</sup> *Ibid.*, p. 609, note 1.






Figure 3. Chemical structure of adenosine triphosphate (ATP).




Figure 4. Chemical structure of adenosine triphosphate chelated with magnesium ion. (MgATP)



g. **Current Flow.** The net result of the many reactions occurring in the mitochondrion is a flow of electrons.<sup>11</sup> This flow is oscillatory in the MgATP crystal.

Gurney determined that removing a negative ion from the interior of a perfect crystal left a vacant lattice point to which a positive charge is associated. If this point is approached by a free electron in the conduction band, the electron will be attracted as from a positive-charged particle. Thus, if the electron loses energy, it may be trapped in the field of the charge. This trapped electron will undergo a series of stationary states toward a series limit. Beyond this limit, a continuum of states will exist. The periodic field of the lattice will cause periodic fluctuations on the wave function of the trapped electron, but the trapped electron will not be prevented from having a definite wave function, in each bound state, which will be like that of an electron within an atom in a vacuum and will be spread over many atomic distances in the crystal.<sup>12</sup>

Similarly, Gurney determined that removing a positive ion from within a perfect crystal left a vacant lattice point to which a negative charge is associated. If this point is approached by a free, positive hole in the filled electronic band, the free, positive hole will be attracted. If the free, positive hole loses energy, it will be trapped in the field of the negative charge. A set of stationary states toward a series limit will be encountered by the trapped, positive hole.<sup>13</sup>

Thus, in either case the current is oscillatory.

h. **Structure of MgATP.** The complete conformational structure of MgATP has not yet been elucidated. A key problem has been the function of the  $Mg^{++}$ . Kothekar *et. al.*<sup>14</sup> made a study of the changes in electronic charge and energy distribution in ATP and ADP after incorporation of  $Mg^{++}$ . They hypothesized that even though  $Mg^{++}$  is not absolutely essential for enzymatic action, its presence facilitates the action by modifying the charge distribution. The most likely position of  $Mg^{++}$  is symmetrical between the three P atoms. Binding between  $Mg^{++}$  and ATP was considered mostly ionic in nature. No detailed, acceptable information is yet available regarding the crystalline structure of MgATP.<sup>15</sup>

4. **Proposed Alternate Mechanism for Energy Production.** There is hardly a material in life organisms which performs only a single function. Therefore, MgATP

<sup>11</sup> Peter C. Hinkle and Richard F. McCarty, "How Cells Make ATP," *Scientific American* 104 (March 1978).

<sup>12</sup> R. W. Gurney, "Ionic Crystals," *Handbook of Physics*, ed. Condon and Odishaw, 2nd edition, 8-49, Section 12 (1967).

<sup>13</sup> *Ibid.*

<sup>14</sup> V. Kothekar, R. S. Tyagi, and R. K. Mishra, "Molecular Orbital Calculations of Magnesium Complexes with ATP and ADP," *Indian J. Biochem and Biophysic*, 10 (4), 279-82 (1973).

<sup>15</sup> O. Kennard, *et. al.*, "The Crystal and Molecular Structure of Adenosine Triphosphate," *Proc. R. Soc. London*, A225, 401-436 (1971).

was examined for an additional and alternate means of energy production which would encompass nuclear fusion reactions in accordance with recognized nuclear theory and practice. The principal method of producing nuclear reactions is by an accelerator – either linear or cyclic. E. O. Lawrence in the 1930s used the cyclotron resonance phenomenon as the basis for the cyclotron particle accelerator. He derived the equations of motion for a charged particle in a uniform, magnetic field with a constant period of revolution so that particles can be accelerated indefinitely in resonance with an oscillatory electric field. Recently, Hunter and McIver developed a small spectroscope based on this principle.<sup>16</sup>

In view of the lack of information regarding the crystalline structure of MgATP or its structure with regard to adjacent molecules of MgATP, it is hypothesized that the molecules of MgATP are situated one above the other, so located that the  $Mg^{++}$  forms a continuous chain. With the exception of the D-ribose unit, the complex or chelate lies in a plane. The D-ribose folds up – a phenomenon called sugar puckering.<sup>17</sup> Since the D-ribose molecules might interfere with each other under these conditions, the MgATP has been rotated so that the D-ribose units would lie  $180^\circ$  apart. A top view of the two molecules is shown in Figure 5. Since the P groups are equidistant from the  $Mg^{++}$ , it follows that they may be  $60^\circ$  apart around a circle whose center is the  $Mg^{++}$ . The acyl oxygen dipoles attached to the P atoms would likewise be situated around the circumference of the circle  $60^\circ$  equidistant. There is evidence that the dipoles might be situated  $30^\circ$  around the circumference in which case four molecules of MgATP would be required for twelve dipoles to be present in the circle.<sup>18</sup> Figures 6 and 7 show top and side views respectively of such an arrangement. The phosphate chain is in the folded configuration and not extended as in anhydrous, inorganic triphosphates. It would form a helix, and torsion angles about the bonds have been calculated.<sup>19</sup> The net effect would be that the  $H^+$  would make a helical path which would be circular as shown in Figure 7.

Dipoles have been found to be energized by unsaturation in the ionic structure – binding with metal ions with each dipole presumably requiring, on the average, one unsaturation or double bond.<sup>20</sup> Hughes and Rideal showed that a dipole is induced in a double bond if it is situated in the  $\alpha$  position but not if it is further away.<sup>21</sup> In contrast, the presence of a double bond in the 4-5 carbon position of

<sup>16</sup> Richard L. Hunter and Robert T. McIver, Jr. "Rapid Scan Ion Cyclotron Resonance Spectroscopy," *American Laboratory*, 9, No. 11, 13 (1978).

<sup>17</sup> D. Kennard, et. al. "The Crystal and Molecular Structure of Adenosine Triphosphate," *Proc. R. Soc. London*, A325, 401-436 (1971).

<sup>18</sup> *Ibid.*

<sup>19</sup> *Ibid.*

<sup>20</sup> O. Dinesh Shah and Jack H. Schulman, *Advanced Chemical Series* 84, 189 (1967).

<sup>21</sup> A. Hughes and E. K. Rideal, *Proc. Roy. Soc. London*, A140, 253 (1933).




Figure 5. Planar view of two molecules of Mg chelated ATP one directly above the other facing each other. The dashed-line structure represents the lower molecule. The  $Mg^{++}$  is at the center of a circle with six dipoles spaced evenly around the circumference - three per molecule of ATP. The adenine part of the adenosine triphosphate straddles the perimeter.




Figure 6. Planar view of four MgATP molecules showing twelve dipoles and only one  $Mg^{++}$  and adenosine molecule. Arrows show  $\alpha$ ,  $\beta$ ,  $\gamma$  rotation of P atoms and, hence, dipoles 1, 2, and 3 when not chelated. Chelation prevents rotation of P atoms and could result in forcing dipoles out of the plane.




Figure 7. Side view of four molecules of MgATP. Possible hydrogen bonding between OH of D-ribose and gamma oxygen is shown.

sphingomyelin increases the surface dipole and, hence, the surface potential of the monolayer.<sup>22</sup> No evidence exists, therefore, for a maximum-distance limitation. In this case, the unsaturation in the adenine molecule comprising four double bonds should, therefore, be sufficient to energize the three dipoles in the phosphate groups. The current formed by the electron flowing in an oscillatory fashion from one  $Mg^{++}$  to another must of necessity be accompanied by an oscillatory magnetic field. This could easily extend out to the perimeter and accelerate the  $H^+$  to relativistic speeds. It would be guided in its path by the dipoles.

An interesting feature noticed in Figure 5 is the location of the adenine rings in the adenosine heterocycles. They sit directly on the circumference or perimeter of the circle formed by the acyl oxygen dipoles and 180° apart from each other. When a conjugated ring molecule is placed in a magnetic field, its electrons are caused to circulate and, in circulating, they generate secondary magnetic fields, i.e., induced magnetic fields. The ring is fully conjugated and, hence, has a rotating ring current only when exposed to a magnetic field.<sup>23</sup> Because of the puckering effect of the D-ribose, the ring slopes and skews downward toward the perimeter. Its induced magnetic field, thus, has a tangential component capable of exerting an accelerating force on the  $H^+$  as it passes by.

Each requirement for a cyclotron particle accelerator has been met on a molecular scale as follows.

a. **Oscillatory Electric Field.** There is a continuous flow of electrons in the mitochondrion. The current oscillates as it flows through the chain of  $Mg^{++}$ s. The electric field accompanying the current oscillates in unison with it.

b. **Presence of Hydrogen Ions.** Hydrogen ions are present in one of the compartments of the mitochondrion.

c. **Auxiliary Control and Propulsion of  $H^+$  by Dipoles.** An  $H^+$  introduced between the components of the sandwich would be attracted by an energized dipole and passed from one dipole to another around a helical circle having a diameter of approximately 30 Angstroms.

d. **Induction of Ring Current in Adenine.** The oscillating electric field induces an oscillating ring current in the adenine ring of the molecule.

e. **Effect of Magnetic Field on Hydrogen Ion.** As the  $H^+$  passes under the ring, it is attracted by the magnetic field accompanying the ring current and accelerated

<sup>22</sup> Q. Dinesh Shah and Jack H. Schulman, *Lipids* 2, 21 (1967).

<sup>23</sup> R. T. Morrison and R. N. Boyd, *Organic Chemistry*, 3rd Edition, 439 (1973).

to a very high speed. The  $H^+$  soon approaches relativistic speeds because the ring current also is probably rotating at such speeds.

f. **Circular Helical Path of Hydrogen Ion.** In a crystal having four molecules of MgATP, the path of the  $H^+$  is a circular helix. As many molecules of MgATP as are necessary would be present to accelerate the  $H^+$  and impart sufficient energy to it to penetrate the nucleus of an atom and convert it to an atom of the next higher number.

### III. DISCUSSION

5. **General.** There were numerous evidences that the premise upon which the study was based was true, i.e., that nuclear transmutations were indeed taking place in life organisms of all types. It was shown that there was a net loss in mass for the transmutations of Na, K, and Mn to Mg, Ca, and Fe respectively. These represented energies in the range of 8.35 MeV to 11.69 MeV. Although there was no information regarding the energy cost in effecting the transmutations, similar transmutations involving hydrogen ions were much less. Thus, there probably was a net gain in energy during the atomic transmutations.

The mitochondrion, well known as the location for energy production in the cell, was closely examined for clues as to the possibility of elemental transmutations also occurring there. The presence of the ion pairs (Na, Mg), (K, Ca), and (Mn, Fe) in the mitochondrion gave strong support to such a conjecture. Moreover, the additional pair (Cu, Zn) found there but not reported by Kervran or others provided added support. These were the only ions reported to have been found in the mitochondrion.

Adenosine triphosphate (ATP) in its chelated form with Mg as MgATP has been pinpointed as the source of energy in the mitochondrion. An examination of its structure indicated that when the molecules are placed one on top of another with the  $Mg^{++}$  in the center, the group resembled a working model of a cyclotron on a molecular scale. When MgATP is taken apart, its components can be burned to also form energy. An analogy would be a windmill or waterwheel made of wood. Both can harness the elements to produce mechanical or electrical energy. When they are taken apart and burned, the components produce thermal energy.

The question of the primary purpose of the transmutations is of considerable interest. From Komaki's work, it would appear that transmutations were used to maintain a balance of certain elements. In the human body, it is evident that health would be considerably impaired if it were attempted to obtain energy continuously from such transmutations. The body can tolerate just so much Na, K, Mg, Ca, etc., and the quantities are small.

No reports have been found of experiments which were made in which energy was measured along with elemental transmutation. Such experiments would be relatively simple to design and perform. The relatively available huge supplies of the elements which have been reported to have been transmuted and the probable large accompanying energy surplus indicate a new source of energy may be in the offing — one whose supply would be unlimited.

#### IV. CONCLUSIONS

6. **Conclusions.** It is concluded that elemental transmutations occurring in life organisms are accompanied by losses in mass representing conversion to thermal energy and that such energy probably is a net gain when compared to the amount required to effect the transmutation.

New Energy Times



## REFERENCES

- Cockcroft, J. and Lewis, B., *Proc. Roy. Soc. A136*, 319; 137, 229 (1932), 154, 246, 26 (1936).
- de Beauregard, O. Costa, *Proc. 3rd Int. Cong. Psych (Tokyo)*, 168 (June 1977).
- Gurney, R. W., "Ionic Crystals," *Handbook of Physics*, Edited by Condon and Odishaw, 2nd Edition, 8-49, Section 12 (1967).
- Hinkle, Peter C. and McCarty, Richard E., "How Cells Make ATP," *Scientific American 104* (March 1978).
- Hughes, A. and Rideal, E. K., *Proc. Roy. Soc. London, A140*, 253 (1933).
- Hunter, Richard L. and McIlver, Robert T., Jr., "Rapid Scan Ion Cyclotron Resonance Spectroscopy," *American Laboratory 19*, No. 11, 13 (1978).
- Jueneman, F. B., "The Transmutation of Species," *Industrial Research, V 19*, No. 13, 11 (Dec 1977).
- Kennard, O. et. al, "The Crystal and Molecular Structure of Adenosine Triphosphate," *Proc. R. Soc. London A325*, 401-436 (1971).
- Kervran, C. L., *Prev. en Biologie de Transmutation a Faible Energy*, Malorinx, Paris (1975).
- Komaki, H., *Rev. de Pathologie Comparee*, 67, 213 (1967); 69, 29 (1969).
- Kothekar, V., Tyagi, R. S., and Mishra, R. K., "Molecular Orbital Calculations of Magnesium Complexes with ATP and ADP," *Indian J. Biochem and Biophysics*, 10(4), 279-82 (1973).
- Makler, Henry R. and Cordes, Eugene H., *Biological Chemistry, VI*, 601, 609, 618 (1966).
- Morrison, R. T. and Boyd, R. N., *Organic Chemistry*, 3rd Edition, 419 (1973).
- Shah, O. Dinesh and Schulman, Jack, *Advanced Chemical Series, 84*, 189 (1967).
- Shah, O. Dinesh and Schulman, Jack, H., *Lipids*, 2, 21 (1967).

**DISTRIBUTION FOR MERADCOM REPORT 2247**

No. Copies	Addressee	No. Copies	Addressee
	<b>Department of Defense</b>	1	Commander US Army Aberdeen Proving Ground ATTN: STEAP-MT-U (GE Branch) Aberdeen Proving Ground, MD 21005
1	Director, Technical Information Defense Advanced Research Projects Agency 1400 Wilson Blvd Arlington, VA 22209		
10	Director Defense Nuclear Agency ATTN: STTL Washington, DC 20305	1	Director US Army Ballistic Research Lab ATTN: DRDAR-TSB-S (STINFO) Aberdeen Proving Ground, MD 21005
12	Defense Documentation Center Cameron Station Alexandria, VA 22314		
	<b>Department of the Army</b>	1	Director US Army Engineer Waterways Experiment Station ATTN: Chief, Library Branch Technical Info Ctr Vicksburg, MS 39180
1	Commander, HQ TRADOC ATTN: ATEN-ME Fort Monroe, VA 23651		
1	HQDA (DAMA-AOA-M) Washington, DC 20310	1	Commander Picatinny Arsenal ATTN: SARPA-TS-S No. 59 Dover, NJ 07801
1	HQDA (DAEN-RDL) Washington, DC 20314	1	Commander US Army Troop Support & Aviation Materiel Readiness Command ATTN: DRSTS-KTE 4300 Goodfellow Blvd St Louis, MO 63120
1	HQDA (DAEN-MCE-D) Washington, DC 20314		
1	Commander US Army Missile Research and Development Command ATTN: DRSMI-RR Redstone Arsenal, AL 35809	1	Commander US Army Electronics Research & Development Command ATTN: DRSEL-GG-TD Fort Monmouth, NJ 07703
1	Director Army Materials and Mechanics Research Center ATTN: DRXMR-STL Technical Library Watertown, MA 02172	1	US Army Aviation School Library P.O. Drawer 0 Fort Rucker, AL 36360
1	US Army Ballistic Research Labs Technical Library DRXBR-LB (Bldg 305) Aberdeen Proving Ground, MD 21005	2	Engineer Representative US Army Standardization Group, UK Box 65, FPO New York 09510
1	Technical Library Chemical Systems Lab Aberdeen Proving Ground, MD 21010	1	Commander Rock Island Arsenal ATTN: SARRI-LPL Rock Island, IL 61201

No. Copies	Addressee	No. Copies	Addressee
1	Learning Resources Center US Army Engineer School Bldg 270 Fort Belvoir, VA 22060	1	Naval Training Equipment Ctr ATTN: Technical Library Orlando, FL 32813
			<b>Department of the Air Force</b>
1	Commander and Director USAFESA ATTN: FESA-RTD Fort Belvoir, VA 22060	1	HQ USAF/RDPS (Mr. Allan Eaffy) Washington, DC 20330
	<b>MERADCOM</b>	1	AFSC/INJ Andrews AFB, MD 20334
1	Commander, DRDME-Z Technical Director, DRDME-ZT Assoc Tech Dir/R&D, DRDME-ZN Assoc Tech Dir/Engrg & Acq, DRDME-ZE Spec Asst/Matl Asmt, DRDME-ZG Spec Asst/Tech Asmt, DRDME-ZK CIRCULATE	1	HQ USAF/PRES ATTN: Mr. Edwin B. Mixon Bolling AFB - Bldg 626 Washington, DC 20332
1	Chief, Crimine Lab, DRDME-N Chief, Engy & Wtr Res Lab, DRDME-G Chief, Elec Pwr Lab, DRDME-E Chief, Cam & Topo Lab, DRDME-R Chief, Mar & Br Lab, DRDME-M Chief, Mech & Constr Eqpt Lab, DRDME-H Chief, Ctr Intrus Lab, DRDME-X Director, Product A&T Directorate, DRDME-T CIRCULATE	1	AFAPL/SFL Wright-Patterson AFB, OH 45433
		1	Department of Transportation Library, FOB 10A, TAD-494.6 800 Independence Ave., SW Washington, DC 20591
			<b>Others</b>
1	Matl Tech Lab, DRDME-V	1	Professor Raymond R. Fox School of Engineering and Applied Science The George Washington Univ Washington, DC 20052
20	Plastics/Ceramics Div, DRDME-VL	20	Secretary of Energy Energy Research & Development Administration 20 Massachusetts Ave NW Washington, DC 20052
3	Tech Reports Ofc, DRDME-WP		
3	Security Ofc (for liaison officers), DRDME-S		
2	Tech Library, DRDME-WC		
1	Plans, Programs & Ops Ofc, DRDME-U		
1	Pub Affairs Ofc, DRDME-I		
1	Ofc of Chief Counsel, DRDME-L		
	<b>Department of the Navy</b>		
1	Director, Physics Program (421) Office of Naval Research Arlington, VA 22217		
1	Director Naval Research Laboratory ATTN: Code 2627 Washington, DC 20375		