## The Puzzle of Excess Heat with No Strong Nuclear Radiation

## Xing Zhong Li

# Dept. of Physics, Tsinghua University Beijing, CHINA

2004,10

## **Answers to Prof. J. Huizenga' Miracles**

- **1. Penetration of Coulomb Barrier**
- 2. No Strong Neutron
- **3. No Strong** γ Radiation



**Compound Nucleus** Decays to

Fastest Channel

(Shortest Lifetime)



**One Dimensional Penetration** 

## **Selective Resonant Tunneling Model**

## **Selective Resonant Tunneling Model**







**1-** Dimensional Penetration

## **Select Damping in Resonance**



#### Phys. Rev. C, 61 (2000),024610



#### Fusion Science & Technology 41(2002) 63

#### **NNDC BNL October 1999**



#### Selective Resonant Tunneling O & NNDC Data +



X. Z. Li, H.Hora, et al., Laser and Particle Beam, 22 No.4 (2004)

**5-Parameter Empirical Fit** 



$$A_1 = 45.95$$
  
 $A_2 = 50200$   
 $A_3 = 1.368 \times 10^{-2}$   
 $A_4 = 1.076$   
 $A_5 = 409$ 

B.H.Duane, "Fusion Cross Section Theory," BWNL-1685,(1972). Naval Research Lab. Plasma Formulary

 $S_0 = e^{i 2 \delta_0}$ 

 $Cot(\delta_0) = W_r + iW_i$ 

 $\sigma_r^{(0)} \approx \frac{\pi}{k^2} (1 - |S_0|^2) \equiv \frac{\pi}{k^2} \left\{ \frac{-4W_i}{W_r^2 + (W_i - 1)^2} \right\}$ 

$$\begin{cases} W_r = 0\\ W_i = O(-1) \end{cases}$$

 $\begin{cases} E = 110 \text{ keV} & \begin{cases} U_{1r} = -47.33 \ MeV \\ \sigma_r^{(0)} = 5.01 \text{ b} & \\ \end{bmatrix} U_{1i} = -115.25 \ keV \end{cases}$ 

 $a = 1.746 \times 10^{-13} (A_1^{1/3} + A_2^{1/3}) cm$ 



Wr



 $\mathcal{T}$ lifelpha



 $\tau_{life} \approx \theta \cdot \tau_{flight} \ (lattice)$  $\tau_{life} \approx \theta^2 \cdot \tau_{flight} \text{ (beam)}$  $\theta^{-2}$  – –*Gamow Factor* 

Czechoslovak Journal of Physics, 49 (1999) 985. Fusion Technology, 36 (1999) 324. Experimental Evidences for Selective Resonant Tunneling

## 1. 3—Deuterons Fusion Reactions

## **2. Tritium Production without n &** $\gamma$

#### Identification of d+d Resonant State



TiD<sub>x</sub> Target  $(5\sim15^{\circ}C, x\sim1.4)$ 

# $d+(d+d) \rightarrow T(4.75 \text{ MeV})+^{3}\text{He}(4.75 \text{ MeV})$ $R(3d) = N_{b}V_{b}\sigma(d+2d)N(2d)$ $N(2d) = R(2d)\tau(2d)$ $\frac{R(3d)}{R(2d)} = N_{b}V_{b}\sigma(d+2d)\tau(2d)$

 $\tau(2d) \approx 10^4 \,\mathrm{sec.} \qquad \tau_{life} \approx \theta \cdot \tau_{flight} \approx 10^{27} \cdot 10^{-23} \,\mathrm{sec.}$ 

## **Tritium Production in D(H)/Pd Systems**

$$d + d \to T + p \qquad \text{(Strong)}$$

$$p + d \to He^{3} + \gamma \qquad \text{(E-M)}$$

$$p + d \to T + e^{+} + v_{e} \qquad \text{(Weak)}$$

$$p + d + e^{-} \to T + v_{e} \qquad \text{(Weak)}$$



#### Selective Resonant Tunneling & NNDC Data

P+d → <sup>3</sup>He + Gamma

*K* – *Capture Lifetime*  $10^{\circ}$  sec  $\beta^+$  – Decay Lifetime 1 sec

 $\theta \cdot \tau_{flight} \approx \tau_{K-capture}$ 

 $10^{25-31} \cdot 10^{-23} \text{ sec} \approx 10^{2-8} \text{ sec}$ 

## **Selectivity of Resonant Tunneling**



P+d→T+e<sup>+</sup>+v<sub>e</sub>  
$$τ_{K-capture}$$
~10<sup>6</sup> sec

$$heta$$
  $au$ flight

 $\theta \tau_{\text{flight}}$ >100 Sec.

## Conclusion

- 1. Excess Heat with No Strong Nuclear Radiation is Feasible. ----- WHY ?
- 2. Deuterium Flux may form a self-sustaining Reactor in Pd.---- HOW ?

Multiple Scattering Theory—Fine Tuning

Fission Reactor—Neutron Diffusion — Escaping Res.

**CMNS Reactor**—**Deuteron Wave**—Lock in Resonance

### **Fine Tuning Mechanism**

• Temperature Variation: 11.8×10<sup>-6</sup> /K

Loading Ratio: D/Pd 0.58→0.78
 Lattice Constant 4.025Å→4.050Å

5×10<sup>-3</sup>/0.2 in 8 ×10<sup>5</sup> seconds